

## CONCERNING CONTINUITY APART FROM A MEAGER SET

JANUŠEK BANIEWSKI AND ZBIGNIEW MOTKOŃCZYK

**ABSTRACT.** Given a  $\sigma$ -ideal  $\mathcal{J}$  of subsets of a space  $X$ , mappings  $f: X \rightarrow Y$  are investigated, such that  $f|X_0$  is continuous for some closed  $X_0 \subset X$  with  $X \setminus X_0 \in \mathcal{J}$ .

**1. Introduction.** Throughout the paper,  $X$  and  $Y$  are topological spaces, and  $\mathcal{J}$  is a  $\sigma$ -ideal of subsets of  $X$ .

The following theorem was first proved by Kanazewski in 1930. (See [K1] and [KM, p. 408]; also [EPR] for a closely related result.)

If  $Y$  is second countable, then the statements

(a)  $f|X_0$  is continuous for some  $X_0 \subset X$  with  $X \setminus X_0 \in \mathcal{J}$  is equivalent to

(b) for every open  $V \subset Y$ ,  $f^{-1}(V) = (U \setminus A) \cup B$ , where  $U$  is open in  $X$ ,  $A \in \mathcal{J}$ , and  $B \in \mathcal{J}$ .

We are interested in characterizing mappings that admit continuous restriction to a set  $X_0 \subset X$  with an extra property in addition to  $X \setminus X_0 \in \mathcal{J}$ . The present paper is to investigate the case of a closed  $X_0$ .

**2. The theorem.** It is an easy observation that if

(i)  $f|X_0$  is continuous for some closed  $X_0 \subset X$  with  $X \setminus X_0 \in \mathcal{J}$ ,

then

(ii) for every open  $V \subset Y$ ,  $f^{-1}(V) = U \setminus A$  with  $U$  open in  $X$  and  $A \in \mathcal{J}$ .

Indeed, let  $X_0$  be as in (i) and take any closed  $F \subset Y$ . For (ii) to hold true,  $f^{-1}(F)$  should have a form of  $E \cup A$  with  $E \subset X$  closed and  $A \in \mathcal{J}$ . Since

$$\begin{aligned} f^{-1}(F) &= [f^{-1}(F) \cap X_0] \cup [f^{-1}(F) \cap (X \setminus X_0)] \\ &= (f|X_0)^{-1}(F) \cup [f^{-1}(F) \cap (X \setminus X_0)], \end{aligned}$$

this is indeed the case.

We are now going to describe circumstances where (i) and (ii) are actually equivalent. To make some statements shorter, let us agree to say that  $f$  is  $\mathcal{J}$ -continuous whenever (i) is satisfied.

**THEOREM.** Let  $Y$  be a regular space and assume that either

- (I)  $Y$  is second countable, or
- (II)  $X$  is hereditarily Lindelöf, or
- (III)  $\mathcal{J}$  consists of all meager sets in  $X$ .

Received by the editors June 14, 1981 and, in revised form, September 24, 1981.

1980 Mathematics Subject Classification (1980 Revision). Primary 54C30; Secondary 26A11.

© 1981 American Mathematical Society  
0002-9947/81 \$1.00 + 1.25 per page

Let  $f: X \rightarrow Y$  be  $\mathcal{A}$ -continuous. Then  $f|X_0$  is continuous for some closed  $X_0 \subset X$  with  $X \setminus X_0 \in \mathcal{A}$ .

**3. Examples.** Before a proof, let us look at two examples showing that the hypotheses of the theorem are to some extent indispensable; namely that

in (I), "second countable" cannot be weakened to "Lindelöf";

in (II), "hereditarily Lindelöf" cannot be weakened to "Lindelöf";

in (III),  $\mathcal{A}$  cannot be just any  $\sigma$ -ideal, and regularity of  $Y$  is essential in all three versions.

**A.** Let  $X = Y = \mathbb{R}$  with topologies as follows. All points except for 0 are isolated in both spaces, while neighborhoods of 0 in  $X$  (respectively, in  $Y$ ) are complements of finite (respectively, countable) sets. These spaces  $X$  and  $Y$  are known to have nice topological properties:  $X$  is compact Hausdorff,  $Y$  is regular Hausdorff and Lindelöf.

Let  $\mathcal{A}$  be the  $\sigma$ -ideal of all countable subsets of  $X$ . Although the identity function  $f: X \rightarrow Y$  is obviously  $\mathcal{A}$ -continuous, its restriction to any closed, co-countable set  $X_0 \subset X$  is discontinuous at 0.

**B.** Now consider  $X = \mathbb{R}$  with its usual topology. Let  $Y = \mathbb{R}$  have a richer topology consisting of all sets  $U \cup A$ , where  $U$  is in the usual topology of  $\mathbb{R}$  and  $A$  is a subset of  $Z = \{1/n: n \in \mathbb{N}\}$ . Thus  $Y$  is Hausdorff and second countable.

Let  $\mathcal{A}$  be the  $\sigma$ -ideal of, say, all meager sets in  $X$  (or all sets of Lebesgue measure 0, or all countable sets). Since  $Z \in \mathcal{A}$ , the identity function  $f: X \rightarrow Y$  is  $\mathcal{A}$ -continuous. On the other hand, the only closed set in  $X$  whose complement is a member of  $\mathcal{A}$  is  $X$  itself. Therefore, the only restriction of  $f$  to be considered is  $f$  itself, and that is not continuous.

**4. Proof of the theorem.** In case (I), let  $\{V_n: n \in \mathbb{N}\}$  be a base for the topology of  $Y$ . By the  $\mathcal{A}$ -continuity of  $f$ , to each  $V_n$  there correspond an open set  $U_n \subset X$  and a set  $A_n \in \mathcal{A}$  so that

$$(1) \quad f^{-1}(V_n) = U_n \setminus A_n.$$

Put

$$(2) \quad J = \bigcup \{A_n: n \in \mathbb{N}\}$$

and

$$(3) \quad X_0 = X \setminus \text{int } J.$$

Of course,  $X_0$  is closed in  $X$  and  $X \setminus X_0 \subset J \in \mathcal{A}$ . To show that  $f|X_0$  is continuous, take any  $x \in X_0$  and any neighborhood  $V$  of  $f(x)$  in  $Y$ . Since  $Y$  is regular, there is an  $n \in \mathbb{N}$  such that  $f(x) \in V_n$  and

$$(4) \quad \text{cl } V_n \subset V.$$

We will show that the corresponding neighborhood  $U_n$  of  $x$  in  $X$ , defined by (1), satisfies  $f(U_n \cap X_0) \subset V$ ; this will prove that  $f|X_0$  is continuous at  $x$ .

Suppose that, on the contrary,  $f(x') \notin V$  for some  $x' \in U_n \cap X_0$ . Then, by (4),  $f(x') \in V \setminus \text{cl } V_n$  and so, by the definition of a base, there is a  $V_m$  with  $f(x') \in V_m$  and

$$(5) \quad V_m \cap V_n = \emptyset.$$

The corresponding set  $U_\alpha$  contains  $x'$ ; therefore

$$(6) \quad x' \in U_\alpha \cap U'_\alpha \cap X_0.$$

On the other hand, (1) and (2) yield  $U_\alpha \subset f^{-1}(V_\alpha) \cup J$  and likewise,  $U'_\alpha \subset f^{-1}(V'_\alpha) \cup J$ . Hence, in view of (4),

$$U_\alpha \cap U'_\alpha \subset f^{-1}(V_\alpha \cap V'_\alpha) \cup J = J$$

and consequently,  $U_\alpha \cap U'_\alpha \subset \text{int } J$ . Now (5) gives  $U_\alpha \cap U'_\alpha \cap X_0 = \emptyset$ , a contradiction to (6).

Cases (II) and (III) can be given a common proof. But first, let us recall a theorem known as Banach Category Theorem. (See [B] and [O, p. 62].)

*If a subset  $X \subset Y$  is locally meager then  $X$  is meager.*

An immediate corollary to the Banach Category Theorem is the following:

*The union of all subsets of  $X$  that are open and meager at the same time is meager.*

In other words, in case (III) of the theorem, the following condition (\*) is satisfied:

(\*) *The union of all members of  $\mathcal{J}$  that are open in  $X$  is a member of  $\mathcal{J}$ .*

The above holds in case (II) as well, since the union of all such sets (in a hereditarily Lindelöf space) coincides with the union of just countably many of them, which is in  $\mathcal{J}$  by the definition of a  $\sigma$ -ideal. Thus (\*) is a property that (II) and (III) have in common.

Let us say that  $f$  is *pointwise  $\mathcal{J}$ -continuous* if, for every  $x \in X$  and for every neighborhood  $V$  of  $f(x)$  in  $Y$ , there exists a neighborhood  $U$  of  $x$  in  $X$  such that  $U \setminus f^{-1}(V) \in \mathcal{J}$ .

Obviously, every  $\mathcal{J}$ -continuous mapping is pointwise  $\mathcal{J}$ -continuous. Therefore, to complete a proof of the theorem, it will suffice to prove the following lemma.

**LEMMA.** *Let  $Y$  be a regular space and let  $f$  be pointwise  $\mathcal{J}$ -continuous. Put*

$$X_0 = X \setminus \bigcup \{G : G \text{ is open in } X \text{ and } G \in \mathcal{J}\}.$$

*Then  $f|X_0$  is continuous.*

**PROOF.** Consider any point  $x \in X_0$  and any neighborhood  $V$  of  $f(x)$  in  $Y$ . We need a neighborhood  $U$  of  $x$  in  $X$  such that  $f(U \cap X_0) \subset V$ .

First, using regularity of  $Y$ , we can select an open set  $W$  containing  $f(x)$  with  $\text{cl } W \subset V$ . Since  $f$  is pointwise  $\mathcal{J}$ -continuous, a required neighborhood  $U$  can now be chosen so that  $U \setminus f^{-1}(W) \in \mathcal{J}$ .

To show that  $f(U \cap X_0) \subset V$ , take any  $x' \in U \cap X_0$ ; since  $\text{cl } W \subset V$ , it will be enough to prove that  $f(x') \in \text{cl } W$ .

Suppose that, on the contrary, there is a neighborhood  $W'$  of  $f(x')$  such that  $W' \cap W = \emptyset$ . By the pointwise  $\mathcal{J}$ -continuity of  $f$ , we have  $U' \setminus f^{-1}(W') \in \mathcal{J}$  for some neighborhood  $U'$  of  $x'$ . Hence

$$U \cap U' = (U \cap U') \setminus f^{-1}(W' \cap W) \subset [U \setminus f^{-1}(W')] \cup [U' \setminus f^{-1}(W)] \in \mathcal{J},$$

and therefore  $U \cap U' \in X \setminus X_0$ , by the definition of  $X_0$ .

We have reached a contradiction, because on the other hand,  $x' \in U \cap U' \cap X_0$ .

5.  *$\mathcal{A}$ -continuity versus pointwise  $\mathcal{A}$ -continuity.* As noted, every  $\mathcal{A}$ -continuous mapping is pointwise  $\mathcal{A}$ -continuous. The converse is not true in general.

For example, let  $X = X_1 \times X_2$ , where  $X_1$  is a discrete uncountable space and  $X_2 = \{0, 1, 1/2, 1/3, \dots\}$  with the usual topology. Let  $\mathcal{A}$  be the  $\alpha$ -ideal of all countable subsets of  $\mathbb{R}$ . Since the space  $X$  is locally countable, any mapping defined on it will automatically be pointwise  $\mathcal{A}$ -continuous. Not necessarily  $\mathcal{A}$ -continuous, though, as witnessed by the characteristic function of the set  $X_1 \times \{0\}$ . Notice that (\*) is not satisfied in this example.

For another example, let  $X$  and  $\mathcal{A}$  be same as above,<sup>1</sup> except that  $X_1$  is now  $\mathbb{R}$  with the usual topology. Let  $Y$  be as in the second example of §3. It is easy to see that the mapping  $f(x_1, x_2) = x_2$  is pointwise  $\mathcal{A}$ -continuous but not  $\mathcal{A}$ -continuous, although (\*) is now satisfied in  $X$ .

However, given both (\*) in  $X$  and regularity of  $Y$ , every pointwise  $\mathcal{A}$ -continuous mapping  $f: X \rightarrow Y$  is  $\mathcal{A}$ -continuous. This follows immediately from the lemmata of §4 and from the observation of §2.

The same is true regardless of regularity of  $Y$ , provided (\*) is replaced with the following stronger condition.

(\*\*\*) For every set  $A \subseteq X$ , if each point  $x \in A$  has a neighborhood  $U(x)$  such that  $U(x) \cap A \in \mathcal{A}$ , then  $A \in \mathcal{A}$ .<sup>2</sup>

More precisely, (\*\*\*) implies that

(T) the family  $\mathcal{F} = \{U \setminus A: U \text{ is open in } X, A \in \mathcal{A}\}$  is closed under arbitrary unions,<sup>3</sup>

which, in turn, implies that every pointwise  $\mathcal{A}$ -continuous mapping defined on  $X$  is  $\mathcal{A}$ -continuous.

Indeed, assume (\*\*\*) and consider any union  $E = \bigcup\{U_i \setminus A_i: i \in S\}$  of members of  $\mathcal{F}$ . Put  $U = \bigcup\{U_i: i \in S\}$  and  $A = U \setminus E$ . Since each  $x \in A$  is in some  $U_i$  and since  $U_i \cap A \subseteq A_i \in \mathcal{A}$ , the condition (\*\*\*) yields  $A \in \mathcal{A}$ , and therefore  $E = U \setminus A \in \mathcal{F}$ . The other implication is equally straightforward.

6. An application. In 1923, Sierpiński and Zygmund [SZ] defined a function  $h: \mathbb{R} \rightarrow \mathbb{R}$  whose restriction to every set of power  $c$  is not continuous. The following is a more general setting of their result. (Compare [KZ, p. 42].)

For every metric separable space  $X$ , and for every complete metric space  $Y$  of power  $c$ , there exists a mapping  $h: X \rightarrow Y$  whose restriction to every set of power  $c$  is not continuous.

The theorem of this paper enables us to further improve that result as follows.

**COROLLARY.** Let  $X$  be a metric separable space and let  $Y$  be a complete metric space. Assume that  $\mathcal{A}$  contains all subsets of  $X$  having power  $< c$ . Then there exists a mapping  $h: X \rightarrow Y$  such that for every set  $X_0 \subseteq X$  with  $X_0 \notin \mathcal{A}$ , the restriction  $f|X_0$  is not  $\mathcal{A}_c$ -continuous,  $\mathcal{A}_c$  being the  $\alpha$ -ideal  $\{A: A \in \mathcal{A}, A \subset X_0\}$ .

<sup>1</sup>Alternatively,  $\mathcal{A}$  can be the  $\alpha$ -ideal of all meager sets in  $X$ .

<sup>2</sup>This condition is still met when either  $\mathcal{F} =$  meager sets (Borel Category Theorem), or  $X$  is a hereditarily Lindelöf space.

<sup>3</sup>And therefore  $\mathcal{F}$  satisfies the axioms for a topology in  $X$ .  $\mathcal{A}$ -continuity then means continuity with respect to that richer topology in  $X$ .

**PROOF.** Let  $h$  be the mapping defined by Sierpiński and Zygmund. Suppose  $h|X_0$  is  $\mathcal{F}_0$ -continuous for some  $X_0 \subset X$  with  $X_0 \notin \mathcal{F}$ . According to our theorem (case (II)), there is a set  $X_1 \subset X_0$  such that  $X_0 \setminus X_1 \in \mathcal{F}_0$  and  $h|X_1$  is continuous. By the choice of  $h$ , this set  $X_1$  has power  $< c$ . Therefore  $X_1 \in \mathcal{F}$ , and consequently  $X_0 = (X_0 \setminus X_1) \cup X_1 \in \mathcal{F}$ . This contradiction ends the proof.

A particular case can be obtained by letting  $\mathcal{F}$  contain all sets of power  $< c$  and no other sets.<sup>4</sup> Then the property of the function  $h$  reads as follows.

*Every set  $X_0$  of power  $c$  contains a point  $x_0$  such that  $h$  takes  $c$ -many points of every neighborhood of  $x_0$  in  $X_0$  outside some fixed neighborhood of  $f(x_0)$ .*

**7. Remarks. A.** In the case when  $\emptyset$  is the only member of  $\mathcal{F}$  that is open in  $X$ , for instance if  $X$  is a Baire space and  $\mathcal{F}$  is the  $\sigma$ -ideal of all meager subsets of  $X$ , condition (i) of §2 coincides with continuity of  $f$ .

Also in that case, our lemma of §4 states that every (pointwise)  $\mathcal{F}$ -continuous mapping  $f: X \rightarrow Y$  is continuous, provided  $Y$  is regular.

**B.** One might be tempted to expect the following two statements to be equivalent as well, at least under some reasonable assumptions on  $X, Y$ , and/or  $\mathcal{F}$ .

(i)  $f|X_0$  is continuous for some open  $X_0 \subset X$  with  $X \setminus X_0 \in \mathcal{F}$ .

(ii) for every open  $V \subset Y$ ,  $f^{-1}(V) = U \cup A$  with  $U$  open in  $X$  and  $A \in \mathcal{F}$ .

As in §2, (i) implies (ii) with no assumptions whatsoever. Unfortunately, the converse is false even when  $X = Y = \mathbb{R}$  (with natural topology) and  $\mathcal{F}$  is any of the standard  $\sigma$ -ideals in  $\mathbb{R}$ . To see this, arrange all rational numbers in a sequence  $q_1, q_2, \dots$  and put

$$f(x) = \begin{cases} 1/n & \text{if } x = q_n, \\ 0 & \text{if } x \text{ is irrational.} \end{cases}$$

Let  $\mathcal{F}$  be any proper  $\sigma$ -ideal containing all rationals. The inverse image of every open set  $V \subset Y$  has a required form, since it is either open or a subset of the rationals. However,  $f$  is discontinuous on every open nonempty set.

#### REFERENCES

- [B] S. Banach, *Réponse au problème de première catégorie*, Fund. Math. 16 (1930), 195–198.  
 [EFZ] A. Engelking, S. Franklin and W. E. Klopka, *On functions having the Baire property*, Bull. Acad. Polon. Sci. Sér. Sci. Math. 17 (1979), 489–491.  
 [E] J. K. Feinstein, *Le problème de Baire dans les espaces métriques*, Fund. Math. 16 (1930), 266–274.  
 [K] ———, *Topology*, Vol. 1, Academic Press, 1966.  
 [KM] E. Kaniowski and A. Mioduski, *Set theory*, 2nd ed., North-Holland, Amsterdam, 1976, Warsaw, 1976.  
 [O] J. C. Oxtoby, *Measure and category*, Springer-Verlag, 1971.  
 [SZ] W. Sierpiński and A. Zygmund, *Sur une fonction qui est discontinue sur tout ensemble de puissance  $\aleph_c$  ou continue*, Fund. Math. 4 (1919), 118–128.

DEPARTMENT OF MATHEMATICS, WARSAW UNIVERSITY, PL 01, 00 WARSAW, POLAND

DEPARTMENT OF MATHEMATICS, YOUNGSDOWN STATE UNIVERSITY, YOUNGSDOWN, OHIO 44155

<sup>4</sup>This  $\mathcal{F}$  is a  $\sigma$ -ideal by König's theorem.